arXiv:1401.4794v1 [math.SP] 20 Jan 2014

On the numerical radii of 2×2 complex matrices

L. Z. Gevorgyan
State Engineering University of Armenia,
Department of Mathematics, E-mail: levgev@hotmail.com

Key words: Numerical radius, projective geometry.
AMS Mathematics Subject Classification 47A12,{Primary}, 51N15, {Secondary}

Abstract

The numerical radius of the general 2×2 complex matrix is calculated.

1 Introduction and preliminaries

Let A be a linear bounded operator, acting in a Hilbert space $(\mathcal{H}, \langle \bullet, \bullet \rangle)$. Denote by $W(A) = \{\langle Ax, x\rangle : \|x\| = 1\}$ the numerical range and by $w(A) = \sup_{\lambda \in W(A)} |\lambda|$ the numerical radius of the operator A. As it is well known, the numerical radius defines a norm on the space of all bounded operators, equivalent to the usual one, i.e.

$$w(A) \leqslant \|A\| \leqslant 2w(A).$$

The importance of the numerical radius is partially motivated by its role in the investigation of many iterative schemes of solution of operator equations Eiermann [1993]. Despite the great abundance of research papers on the numerical range and the numerical radius, the exact calculations of the numerical radius of general non-normal operator are very scarce. A few known

results concern the $n-$ dimensional Jordan block

$$J_n = \begin{pmatrix} 0 & 1 & 0 & \cdots & 0 \\ 0 & 0 & 1 & \cdots & 0 \\ \cdot & \cdot & \cdot & \cdot & \cdot \\ 0 & 0 & 0 & \cdots & 1 \\ 0 & 0 & 0 & \cdots & 0 \end{pmatrix}$$

$(w(J_n) = \cos\frac{\pi}{n+1})$ and the Volterra integration operator

$$(Vf)(x) = \int_0^x f(t)dt, f \in L^2(0;1)$$

$(w(V) = 1/2)$.

In the sequel we consider the simplest case of two-dimensional space.

2 Main result

As any operator in a finite-dimensional space may be reduced to the Schur's upper triangular form, we get the matrix

$$A = \begin{pmatrix} \lambda_1 & \lambda_3 \\ 0 & \lambda_2 \end{pmatrix}.$$

Recall that the numerical range of an operator, acting in a two-dimensional space is an elliptical disk (Halmos [1982], ch. 22) having as foci two eigenvalues $\{\lambda_1, \lambda_2\}$ and

$$\left\{tr(A^*A) - |\lambda_1|^2 - |\lambda_2|^2\right\}^{1/2}$$

as minor axis $2b$.

We have $b = |\lambda_3|/2$, the distance between foci is $2c = |\lambda_2 - \lambda_1|$, hence major axis is $2a = \sqrt{|\lambda_2 - \lambda_1|^2 + |\lambda_3|^2}$. The center of symmetry of the ellipse has the affix $m = \frac{\lambda_2+\lambda_1}{2}$ and whole ellipse is rotated by the angle $\varphi = \arg(\lambda_2 - \lambda_1)$. The equation of the ellipse is

$$\frac{((x - \operatorname{Re} m)\cos\phi + (y - \operatorname{Im} m)\sin\phi)^2}{a^2} + \frac{((x - \operatorname{Re} m)\sin\phi - (y - \operatorname{Im} m)\cos\phi)^2}{b^2} - 1 = 0.$$

After elementary transformations it may be reduced to the form

$$G(x,y) = Ax^2 + 2Bxy + Cy^2 + 2Dx + 2Ey + F = 0,$$

where

$$A = \frac{\cos^2\phi}{a^2} + \frac{\sin^2\phi}{b^2}, B = \sin\phi\cos\phi\left(\frac{1}{a^2} - \frac{1}{b^2}\right), C = \frac{\sin^2\phi}{a^2} + \frac{\cos^2\phi}{b^2},$$
$$D = -A\operatorname{Re} m - B\operatorname{Im} m, E = -B\operatorname{Re} m - C\operatorname{Im} m,$$
$$F = \left(\frac{\cos\phi\operatorname{Re} m + \sin\phi\operatorname{Im} m}{a}\right)^2 + \left(\frac{\cos\phi\operatorname{Im} m - \sin\phi\operatorname{Re} m}{b}\right)^2 - 1.$$

The square of numerical radius $w(A)$ may be found as the greatest value of the sum x^2+y^2, where x and y are constrained by the equation $G(x,y) = 0$. According to the Lagrange multiplier method, the auxiliary function

$$L(x,y,\lambda) = x^2 + y^2 + \lambda G(x,y)$$

should be introduced and the system of equations

$$\begin{cases} \frac{\partial L}{\partial x} = 0, \\ \frac{\partial L}{\partial y} = 0, \\ \frac{\partial L}{\partial \lambda} = 0 \end{cases}$$

will be solved.

We prefer another approach. The equation of the circumference is

$$\Gamma(x,y) = x^2 + y^2 - R^2 = 0.$$

If the ellipse lies inside the circle and touches the circumference, then they have a common tangent, therefore

$$\frac{Ax + By + D}{Bx + Cy + E} = \frac{x}{y}.$$

Finally, we get the set of simultaneous equations

$$\begin{cases} Ax^2 + 2Bxy + Cy^2 + 2Dx + 2Ey + F = 0, \\ Bx^2 + (C - A)xy - By^2 + Ex - Dy = 0. \end{cases} \quad (1)$$

Note that the second curve is a hyperbola. The solution of this set may be reduced to a quartic polynomial equation. Using some notions of projective geometry (see Richter-Gebert [2011], ch. 11), it is possible to avoid cumbersome calculations. Introducing the homogeneous coordinates $\begin{pmatrix} x & y & 1 \end{pmatrix}^t$ of the point $(x;y)$, two matrices

$$G = \begin{pmatrix} A & B & D \\ B & C & E \\ D & E & F \end{pmatrix} \text{ and } H = \frac{1}{2}\begin{pmatrix} 2B & C-A & E \\ C-A & -2B & -D \\ E & -D & 0 \end{pmatrix}$$

may be associated with (1). Two conics define a pencil of conics $\alpha H + \beta G; \alpha, \beta \in \mathbb{R}$, passing through the intersection points of G and H. Solving the generalized eigenvalue problem $\det(H - \lambda G) = 0$ we get the degenerated conics $C = H - \lambda_0 G, \det(C) = 0$. Now we calculate the adjugate matrix $J = Ad(C)$. Maybe the simplest way is the use of the formula $J = -(a_1 I + a_2 C + C^2)$, where $a_1 = \frac{1}{2}(tr^2(C) - tr(C^2)), a_2 = -tr(C)$. Denote $p_1 = \sqrt{J_{11}}, p_2 = \sqrt{J_{22}}, p_3 = \sqrt{J_{33}}$, where the signs before the radicals are chosen in a such way that $\{p_1, p_2, p_3\}^T \cdot \{p_1, p_2, p_3\} = J$ and construct the antisymmetric matrix

$$P = \begin{pmatrix} 0 & p_3 & -p_2 \\ -p_3 & 0 & p_1 \\ p_2 & -p_1 & 0 \end{pmatrix}.$$

Now we choose the parameter α to satisfy the condition $K = C - \alpha P, rank(K) = 1$. Any row (or column) of K defines a degenerate conics (a straight line) l. Replacing the first equation in (1) by $l = 0$ we may found easily (solving a quadratic equation) the coordinates of the intersection points. One of them will supply the minimum of $\sqrt{x^2 + y^2}$ and the second point is at the distance $w(A)$ from the origin of the coordinate system.

We append below a MatLab program, which admits as input a 2×2 upper triangular matrix and calculates its numerical radius.

3 Appendix

```
function numradius(A)
 nor=norm(A);
 b=.5*abs(A(1,2));
```

```
c=.5*abs(A(1,1)-A(2,2));
a=sqrt(b^ 2+c^ 2);
m=.5*(A(1,1)+A(2,2));
k=real(m);l=imag(m);
phi=angle(A(2,2)-A(1,1));
s=linspace(0,2*pi,2000);
u=a*cos(s);v=b*sin(s);
R=[cos(phi) -sin(phi);sin(phi) cos(phi)];
K=R*[u;v];
z=K(1,:)+k+i*(K(2,:)+l);
plot(real(z),imag(z),'r')
axis 'equal'
A=cos(phi)^ 2/a^ 2+sin(phi)^ 2/b^ 2;
B=sin(phi)* cos(phi)*(1/a^ 2-1/b^ 2);
C=cos(phi)^ 2/b^ 2+sin(phi)^ 2/a^ 2;
D=-k*A-l*B;
E=-k*B-l*C;
F=(k*cos(phi)+l*sin(phi))^ 2/a^ 2+(l*cos(phi)-k*sin(phi))^ 2/b^ 2-1;
G=[A B D;B C E;D E F];
H=.5*[2*B C-A E;C-A -2*B -D;E -D 0];
e=eig(H,G);
for j=1:3
if isreal(e(j))
c=e(1);
end
end
C=H-c*G;
s1=trace(C);s2=trace(C^ 2);a2=-s1;a1=-.5*(s1*a2+s2);
Ad=-(a1*eye(3)+a2*C+C^ 2);
p(1)=sqrt(Ad(1,1));
p(2)=sign(Ad(1,2))*sqrt(Ad(2,2));
p(3)=sign(Ad(1,3))*sqrt(Ad(3,3));
P=[0 p(3) -p(2);-p(3) 0 p(1);p(2) -p(1) 0];
e=eig(C,P);
e(4)=sqrt((C(1,2)^ 2-C(1,1)*C(2,2)))/p(3);
e(5)=sqrt((C(3,2)^ 2-C(3,3)*C(2,2)))/p(1);
e(6)=sqrt((C(1,3)^ 2-C(1,1)*C(3,3)))/p(2);
for j=1:6
```

```
    K=C-e(j)*P;
    r(j)=rank(K);
    l=mean(K(:,1));
    n=mean(K(:,2));
    q=mean(K(:,3));
    [x,y]=solve('H(1,1)*x^ 2+H(2,2)*y^ 2+2*H(1,2)*x*y+...
2*H(1,3)*x+2*H(2,3)*y=0','l*x+n*y+q=0');
    m1=eval(sqrt(x(1)^ 2+y(1)^ 2));
    m2=eval(sqrt(x(2)^ 2+y(2)^ 2));
    w(j)=max(m1,m2);
    end
    for j=1:6
    if isreal(w(j))&& w(j)≤ nor&&(r(j)==1)
    numr=w(j);
    end
    end
```

References

Michael Eiermann. Fields of values and iterative methods. *Linear Algebra Appl.*, 180:167–197, 1993. ISSN 0024-3795. doi: 10.1016/0024-3795(93)90530-2. URL http://dx.doi.org/10.1016/0024-3795(93)90530-2.

Paul Richard Halmos. *A Hilbert space problem book*, volume 19 of *Graduate Texts in Mathematics*. Springer-Verlag, New York, second edition, 1982. ISBN 0-387-90685-1. Encyclopedia of Mathematics and its Applications, 17.

Jürgen Richter-Gebert. *Perspectives on projective geometry*. Springer, Heidelberg, 2011. ISBN 978-3-642-17285-4. doi: 10.1007/978-3-642-17286-1. URL http://dx.doi.org/10.1007/978-3-642-17286-1. A guided tour through real and complex geometry.